arXiv:math/0702868v1 [math.NT] 28 Feb 2007

A NOTE ON p-ADIC q-EULER MEASURE

Hacer OZDEN[*], **Yilmaz SIMSEK**[**], **Seog-Hoon RIM**[***], **Ismail Naci CANGUL**[*]

[*] *University of Uludag, Faculty of Arts and Science, Department of Mathematics, 16059 Bursa*, Turkey
[**] *University of Akdeniz, Faculty of Arts and Science, Department of Mathematics, 07058 Antalya*, Turkey

[***] *University of Kyungpook, Department of Mathematical Education, Taegu, 702-701, S. KOREA*

E-Mail: hozden@uludag.edu.tr, simsek@akdeniz.edu.tr, shrim@knu.ac.kr and cangul@uludag.edu.tr

Abstract

In this paper, we will investigate some interesting properties of the modified q-Euler numbers and polynomials. The main purpose of this paper is to construct p-adic q-Euler measure on $\mathbb{Z}_p$.

2000 Mathematics Subject Classification. Primary 11S40,11S80; Secondary 11B68.
Key Words and Phrases. *p-adic q-integral, Euler numbers, Euler polynomials, p-adic Volkenborn integral, q-Euler measure*

1. Introduction, Definitions and Notations

Let p be a fixed odd prime. Throughout this paper $\mathbb{Z}_p$, $\mathbb{Q}_p$, $\mathbb{C}$ and $\mathbb{C}_p$ will respectively denote the ring of p-adic rational integers, the field of p-adic rational numbers, the complex number field and the completion of the algebraic closure of $\mathbb{Q}_p$. Let v_p be the normalized exponential valuation of $\mathbb{C}_p$ with $|p|_p = p^{-v_p(p)} = 1/p$, see [1], [2], [3]. When one talks of q-extensions, q is variously considered as an indeterminate, a complex $q \in \mathbb{C}$, or a p-adic number $q \in \mathbb{C}_p$. If $q \in \mathbb{C}$, one normally assumes $|q| < 1$. If $q \in \mathbb{C}_p$, then we assume $|q-1| < p^{-\frac{1}{p-1}}$, so that $q^x = \exp(x \log q)$ for $|x|_p \leq 1$, see [4], [5], [6]. It was well known that Euler numbers are defined by

$$\frac{2}{e^t+1} = \sum_{n=0}^{\infty} E_n \frac{t^n}{n!}, \text{ see [1], [2], [3], [4], [5], [6].}$$

In the recent paper [3], the q-extension of Euler numbers are defined inductively by

$$E_{0,q}=1,\ q(qE+1)^n+E_{n,q}=\begin{cases}[2]_q & if\ n=0\\ 0 & if\ n\neq 0\end{cases}$$

with the usual convention of replacing E^m by $E_{m,q}$. In [1], the definition of modified q-Euler numbers $\mathcal{E}_{0,q}$ is introduced by

$$\mathcal{E}_{0,q}=\frac{[2]_q}{2},\ (q\mathcal{E}+1)^k-\mathcal{E}_{k,q}=\begin{cases}[2]_q & if\ k=0\\ 0 & if\ k>0\end{cases}$$

with the usual convention of replacing $\mathcal{E}^i$ by $\mathcal{E}_{i,q}$.

For a fixed positive integer d with $(p,d)=1$, set

$$X_d=\lim_{\overleftarrow{N}}\mathbb{Z}/dp^N\mathbb{Z},\quad X_1=\mathbb{Z}_p,$$

$$X^*=\underset{\substack{0<a<dp\\(a,p)=1}}{\cup}(a+dp\mathbb{Z}_p),$$

$$a+dp^N\mathbb{Z}_p=\left\{x\in\ X\ :\ x\equiv a\ (\mathrm{mod}\ \ dp^N)\right\},$$

where $a\in\mathbb{Z}$ satisfies the condition $0\leq a<dp^N$, see [6].

We say that f is a uniformly differentiable function at a point $a\in\mathbb{Z}_p$, and write $f\in UD(\mathbb{Z}_p)$, if the difference quotient

$$F_f(x,y)=\frac{f(x)-f(y)}{x-y}$$

has a limit $f'(a)$ as $(x,y)\to(a,a)$. For $f\in UD(\mathbb{Z}_p)$, an invariant p-adic q-integral was defined by

$$I_q(f)=\int_{\mathbb{Z}_p}f(x)d\mu_q(x)=\lim_{N\to\infty}\frac{1}{[p^N]_q}\sum_{x=0}^{p^N-1}f(x)g^x,\ \text{see [8]}.$$

The q-extension of $n\in\mathbb{N}$ is defined by

$$[n]_q=\frac{1-q^n}{1-q}=1+q+q^2+...+q^{n-1},$$

and

$$[n]_{-q}=\frac{1-(-q)^n}{1-(-q)}=1-q+q^2-...+(-q)^{n-1},\ \text{see [3],[4],[5]}.$$

The modified p-adic q-integral on $\mathbb{Z}_p$ is defined by

$$I_{-q}(f)=\int_{\mathbb{Z}_p} f(x)d\mu_{-q}(x),$$

where $d\mu_{-q}(x)=\lim_{q\to -q} d\mu_q(x)$. In this paper, we will investigate some interesting properties of the modified q-Euler numbers and polynomials. The purpose of this paper is to construct p-adic q-Euler measure on $\mathbb{Z}_p$.

2. p-adic q-Euler Measure on $\mathbb{Z}_p$

q-Euler numbers are known by

$$\mathcal{E}_{0,q}=\frac{[2]_q}{2},\ (q\mathcal{E}+1)^n-\mathcal{E}_{n,q}=\begin{cases}[2]_q & if\ n=0\\ 0 & if\ n>0\end{cases}$$

with the usual convention of replacing $\mathcal{E}^i$ by $\mathcal{E}_{i,q}$, see [1]. It was known that the q-Euler numbers can be represented by p-adic q-integrals on $\mathbb{Z}_p$ as follows:

$$\mathcal{E}_{n,q}=\int_{\mathbb{Z}_p} q^{-x}[x]_q^n d\mu_{-q}(x)=[2]_q\left(\frac{1}{1-q}\right)^n\sum_{l=0}^{n}(-1)^l\binom{n}{l}\frac{1}{1+q^l},\text{ see [1].}$$

We now also consider the q-Euler polynomials as follows:

$$\mathcal{E}_{n,q}(x)=\int_{\mathbb{Z}_p} q^{-x}[t+x]_q^n d\mu_{-q}(t)=\sum_{l=0}^{n}\binom{n}{l}q^{xl}\mathcal{E}_{l,q}[x]_q^{n-l}.$$

Thus, we note that

$$\mathcal{E}_{n,q}(x)=[d]_q^n\frac{[2]_q}{[2]_{q^d}}\sum_{a=0}^{d-1}(-1)^a\,\mathcal{E}_{n,q^d}\left(\frac{n+a}{d}\right),\text{ see [1].}\tag{2.1}$$

Let χ be the Dirichlet's character with odd conductor $d\in\mathbb{N}$, and let $F_{\chi,q}(t)$ be the generating function of $\mathcal{E}_{n,\chi,q}$ as follows:

$$F_{\chi,q}(t)=[2]_q\sum_{n=0}^{\infty}(-1)^n\chi(n)e^{[n]}q^t=\sum_{n=0}^{\infty}\mathcal{E}_{n,\chi,q}\frac{t^n}{n!},\text{ see [1].}\tag{2.2}$$

From (2.2) we derive

$$\mathcal{E}_{n,\chi,q}=[d]_q^n\frac{[2]_q}{[2]_{q^d}}\sum_{a=0}^{d-1}(-1)^a\chi(a)\mathcal{E}_{n,q^d}\left(\frac{a}{d}\right).\tag{2.3}$$

For any positive integers N, k and d (odd), let $\mu_k^*=\mu_{k,q;E}^*$ be defined by

$$\mu_k^*(a+dp^N\mathbb{Z}_p)=(-1)^a\,[dp^N]_q^k\frac{[2]_q}{[2]_{q^{dp^N}}}\mathcal{E}_{k,q^{dp^N}}\left(\frac{a}{dp^N}\right).\tag{2.4}$$

Then we see that

$$\begin{aligned}
&\sum_{i=0}^{p-1}\mu_k^*(a+idp^N+dp^{N+1}\mathbb{Z}_p)\\
=\;& [dp^{N+1}]_q^k\frac{[2]_q}{[2]_{q^{dp^{N+1}}}}\sum_{i=0}^{p-1}(-1)^{a+idp^N}\,\mathcal{E}_{k,q^{dp^{N+1}}}\left(\frac{a+idp^N}{dp^{N+1}}\right)\\
=\;& (-1)^a\,[dp^{N+1}]_q^k\frac{[2]_q}{[2]_{q^{dp^{N+1}}}}\sum_{i=0}^{p-1}(-1)^i\,\mathcal{E}_{k,(q^{dp^N})^p}\left(\frac{\frac{a}{dp^N}+i}{p}\right)\\
=\;& (-1)^a\,[dp^N]_q^k\frac{[2]_q}{[2]_{q^{dp^N}}}\left(\frac{[2]_{q^{dp^N}}}{[2]_{(q^{dp^N})^p}}[p]^k_{q^{dp^N}}\sum_{i=0}^{p-1}(-1)^a\,\mathcal{E}_{k,(q^{dp^N})^p}\left(\frac{\frac{a}{dp^N}+i}{p}\right)\right)\\
=\;& (-1)^a\,[dp^N]_q^k\frac{[2]_q}{[2]_{q^{dp^N}}}\mathcal{E}_{k,q^{dp^N}}\left(\frac{a}{dp^N}\right)\\
=\;& \mu_k^*(a+dp^N\mathbb{Z}_p).
\end{aligned}$$

It is easy to see that $|\mu_k^*| \;\leq\; M$, for some constant M. Therefore we obtain the following:

Theorem 1. *For any positive integers* $N,\ k$ *and* d *(odd), let* $\mu_k^*=\mu_{k,q;E}^*$ *be defined by*

$$\mu_k^*(a+dp^N\mathbb{Z}_p)=.\,(-1)^a\,[dp^N]_q^k\frac{[2]_q}{[2]_{q^{dp^N}}}\varepsilon_{k,q^{dp^N}}\left(\frac{a}{dp^N}\right).$$

Then μ_k^* *is a measure on* X.

From the definition of μ_k^*, we derive the following:

$$\begin{aligned}
&\int_X\chi(x)d\mu_k^*(x)=\lim_{N\to\infty}\sum_{a=0}^{dp^N-1}\chi(a)\mu_k^*(a+dp^N\mathbb{Z}_p)\\
&=\lim_{N\to\infty}\sum_{x=0}^{dp^N-1}(-1)^x\chi(x)[dp^N]_q^k\frac{[2]_q}{[2]_{q^{dp^N}}}\mathcal{E}_{k,q^{dp^N}}\left(\frac{x}{dp^N}\right)\\
&=\lim_{N\to\infty}[d]_q^k[p^N]^k_{q^d}\frac{[2]_q}{[2]_{q^d}}\frac{[2]_{q^d}}{[2]_{(q^d)^{p^N}}}\sum_{x=0}^{dp^N-1}(-1)^x\chi(x)\mathcal{E}_{k,q^{dp^N}}\left(\frac{x}{dp^N}\right)
\end{aligned}\tag{2.5}$$

$$\begin{aligned}
=\;& [d]_q^k\frac{[2]_q}{[2]_{q^d}}\lim_{N\to\infty}\frac{[2]_{q^d}}{[2]_{(q^d)^{p^N}}}[p^N]^k_{q^d}\sum_{a=0}^{d-1}\sum_{x=0}^{p^N-1}(-1)^{a+dx}\chi(a+dx)\mathcal{E}_{k,(q^d)^{p^N}}\left(\frac{\frac{a}{d}+x}{p^N}\right)\\
=\;& [d]_q^k\frac{[2]_q}{[2]_{q^d}}\lim_{N\to\infty}\left\{\sum_{a=0}^{d-1}\frac{[2]_{q^d}}{[2]_{(q^d)^{p^N}}}[p^N]^k_{q^d}\sum_{x=0}^{p^N-1}(-1)^a\chi(a)\mathcal{E}_{k,(q^d)^{p^N}}\left(\frac{\frac{a}{d}+x}{p^N}\right)\right\}
\end{aligned}$$

and by equation (2.1), this becomes

$$\int_X \chi(x)d\mu^*_k(x) = [d]_q^k \frac{[2]_q}{[2]_{q^d}} \sum_{a=0}^{d-1} (-1)^a \chi(a) \mathcal{E}_{n,q^d}\left(\frac{a}{d}\right).$$

By using (2.3) and (2.5), we obtain the following:

Theorem 2. *For any positive integer k, we have*

$$\int_X \chi(x)d\mu^*_k(x) = \mathcal{E}_{k,\chi,q}.$$

From Theorem 1 and equation (2.1), we note that

$$\begin{aligned}
&\mu^*_k(a+dp^N\mathbb{Z}_p) \\
= \;& (-1)^a \frac{[2]_q}{[2]_{q^{dp^N}}} [dp^N]_q^k \mathcal{E}_{k,q^{dp^N}}\left(\frac{a}{dp^N}\right) \\
= \;& \left(\sum_{l=0}^{k} \left(\begin{array}{c} k \\ l \end{array}\right) q^{dp^N \frac{a}{dp^N} l} \mathcal{E}_{l,q^{dp^N}} [\frac{a}{dp^N}]_{qdp^N}^{k-l}\right) (-1)^a \frac{[2]_q}{[2]_{q^{dp^N}}} [dp^N]_q^k \\
= \;& \mathcal{E}_{0,q} [\frac{a}{dp^N}]_{qdp^N}^k (-1)^a \frac{[2]_q}{[2]_{q^{dp^N}}} [dp^N]_q^k + (-1)^a \frac{[2]_q}{[2]_{q^{dp^N}}} [dp^N]_q^k \sum_{l=1}^{k} \left(\begin{array}{c} k \\ l \end{array}\right) q^l \\
= \;& \mathcal{E}_{0,q^{dp^N}} [\frac{a}{dp^N}]_{qdp^N}^k (-1)^a \frac{[2]_q}{[2]_{q^{dp^N}}} [dp^N]_q^k + (-1)^a \frac{[2]_q}{[2]_{q^{dp^N}}} [dp^N]_q^k \cdot \\
& \sum_{l=1}^{k} \left(\begin{array}{c} k \\ l \end{array}\right) q^l \mathcal{E}_{l,q^{dp^N}} \left[\frac{a}{dp^N}\right]_{q^{dp^N}}^{k-l} \\
= \;& \frac{[2]_{q^{dp^n}}}{2} \frac{[a]_q^k}{[dp^N]_q^k} (-1)^a \frac{[2]_q}{[2]_{q^{dp^N}}} [dp^N]_q^k + (-1)^a \frac{[2]_q}{[2]_{q^{dp^N}}} [dp^N]_q^k \cdot . \\
& \sum_{l=1}^{k} \left(\begin{array}{c} k \\ l \end{array}\right) q^l \mathcal{E}_{l,q^{dp^N}} \left(\frac{[a]_q}{[dp^N]_q}\right)^{k-l}.
\end{aligned}$$

Thus, we have

$$\begin{aligned}
\lim_{N\to\infty} \mu^*_k(a+dp^N\mathbb{Z}_p) &= \frac{[2]_q}{2}(-1)^a [a]_q^k \\
&= q^a \frac{[2]_q}{2}(-1)^a q^{-a} [a]_q^k \\
&= q^{-a}[a]_q^k \lim_{N\to\infty} \mu_{-q}(a+dp^N\mathbb{Z}_p),
\end{aligned}$$

where d is a positive odd integer. Therefore we obtain the following:

Theorem 3. *For any positive integer k, we have*

$$q^{-x}[x]_q^k d\mu_{-q}(x) = d\mu_k^*(x).$$

Corollary 1. *Let k be a positive integer. Then we have*

$$\mathcal{E}_{k,\chi,q} = \int_X \chi(x) d\mu_k^*(x) = \int_X \chi(x) q^{-x}[x]_q^k d\mu_{-q}(x).$$

Let $\overline{d} = (d, p)$ be the least common multiple of the conductor d of X and p, and let $\mathcal{E}_{n,\chi,q}$ denote the n-th generalized q-Euler number belonging to the character χ. Then we have the q-analogue form of Witt's formula in the cyclotomic field $\mathbb{Q}_p(\chi)$ as follows:

For all $n \geq 0$, we have

$$\mathcal{E}_{n,\chi,q} = \lim_{\rho\to\infty} \frac{[2]_q}{2} \sum_{x=1}^{\overline{d}p^\rho} (-1)^{\chi} \chi(x)[x]_q^n. \tag{2.6}$$

Herein as usual we set $\chi(x) = 0$ if x is not prime to the conductor d. From (2.6) we derive

$$\begin{aligned}
\mathcal{E}_{n,\chi,q} &= \lim_{\rho\to\infty} \frac{[2]_q}{2} \sum_{1\leq x\leq \overline{d}p^\rho}^{*} (-1)^x \chi(x)[x]_q^n \\
&\qquad + \lim_{\rho\to\infty} \frac{[2]_q}{[2]_{q^p}} \frac{[2]_{q^p}}{2} \sum_{1\leq x\leq \overline{d}p^\rho}^{*} (-1)^x \chi(p)\chi(x)[p]_q^n [x]_{q^p}^n \\
&= \lim_{\rho\to\infty} \frac{[2]_q}{2} \sum_{1\leq x\leq \overline{d}p^\rho}^{*} (-1)^x \chi(x)[x]_q^n \\
&\qquad + \frac{[2]_q}{[2]_{q^p}} \lim_{\rho\to\infty} \frac{[2]_{q^p}}{2} \chi(p)[p]_q^n \sum_{1\leq x\leq \overline{d}p^\rho}^{*} (-1)^x \chi(x)[x]_{q^p}^n,
\end{aligned}$$

where $*$means taking the sum over the rational integers prime to p in the given range. Thus we have

$$\mathcal{E}_{n,\chi,q} = \lim_{\rho\to\infty} \frac{[2]_q}{2} \sum_{1\leq x\leq \overline{d}p^\rho}^{*} (-1)^{\chi} \chi(x)[x]_q^n + \frac{[2]_q}{[2]_{q^p}} \chi(p)[p]_q^n \, \mathcal{E}_{n,\chi,q^p},$$

that is,

$$\begin{aligned}\mathcal{E}_{n,\chi,q}-\frac{[2]_q}{[2]_{q^p}}\chi(p)[p]_q^n\ \mathcal{E}_{n,\chi,q^p} &= \lim_{\rho\to\infty}\frac{[2]_q}{2}\sum_{1\le x\le \overline{d}p^{\rho}}^{*}(-1)^{\chi}\,\chi(x)[x]_q^n \\ &= \int\limits_{X^*}(-1)^{x}\chi w^n(x)\left\langle x\right\rangle_q^n\ d\mu_{-q}(x),\end{aligned}$$

where $\left\langle x\right\rangle_q=\frac{[x]_q}{w(x)}$, and $w(x)$ is the Teichmüller character.

Acknowledgement 1. *The first and fourth authors are supported by the research fund of Uludag University project no: F-2006/40. The second author is supported by the research fund of Akdeniz University.*

REFERENCES

[1] Kim, T., *The modified q-Euler numbers and polynomials*, arxivmath.NT/0702523.
[2] Kim, T., *On a q-analogue of the p-adic log gamma functions*, J. Number Theory, 16 (1999), 320-329.
[3] Kim, T., *On the q-extension of Euler and Genocchi numbers,* J. Math. Anal. Appl., 326 (2001),1458-1465.
[4] Kim, T., *Sums of powers of consecutive q-integers,* Advan. Stud. Contemp. Math., 9 (2004), 15-18.
[5] Kim, T., *An invariant p-adic q-integral on $\mathbb{Z}_p$*, Appl. Math. Letters (in press).
[6] Srivastava, H. M., Kim, T., Simsek, Y., *q-Bernoulli numbers and polynomials associated with multiple q-zeta functionsand basic L-series,* Russ. J. Math. Phts. 12 (2005), 241-268.
[7] Kim, T., *On p-adic q-Bernoulli numbers,* J. Korean Math. Soc., 39 (2000), 21-30.
[8] Kim, T., *A new approach to q-zeta function,* J. Comput. Analy. Appl., 9 (2007), 395-400.